

post procedure instructions | GASTROSCOPY

Your procedure involved sedation and the following applies to you:

- **Be** aware of the risk of falling and take the appropriate measures;
- You should not drive a vehicle for 24 hours after your procedure;
- ❖ You should avoid alcohol for 24 hours after your procedure;
- You may be dehydrated and will need to increase your fluid intake for the next 24-hours.

Other things to be aware of

- You may return to a normal diet and fluids
- Remove the IV pressure pad after 3 hours.
- If any signs of infection appear at the cannula site please contact our office and make an appointment to see your GP.

You may experience some discomfort over the next 24-48 hours Some of these symptoms may be:

- Discomfort in the chest area from biopsy sites, wind pain and/or a sore throat
- You may pass a small amount of blood in your mucous when coughing or blowing your nose.

These symptoms usually settle without treatment, however if you have persistent pain, bleeding or develop a temperature you should seek medical assistance.

Please contact the facility if you experience any signs or symptoms of infection within 30 days that could be related to your procedure.

Please contact your specialist or general physician during office hours or your nearest medical facility after hours.

Gastrotract 02 6234 7900

Brindabella Gastroenterology 02 6281 4827 Canberra Gastroenterology 02 6162 1515

Dr PK Tan 02 6156 0291

Dr Ian Davis, Dr Ram Ganesalingam, Dr Paul Pavli 02 6281 0455

Dr Jeff van Gangelen 02 6492 1155

A Member of **CUFA**day hospitals group

